arXiv:1203.2913v1 [math.AG] 13 Mar 2012

ON THE TATE CONJECTURE FOR THE FANO SURFACES OF CUBIC THREEFOLDS

XAVIER ROULLEAU

ABSTRACT. The Fano surface of a smooth cubic threefold $F \hookrightarrow \mathbb{P}^4$ parametrizes the lines on F. We prove that a Fano surface verify the Tate conjecture over finite fields and number fields. We give an algorithm to compute the zeta function of the intermediate Jacobian of the cubic when the field of definition is finite and we compute the zeta function of various examples of Fano surfaces. We obtain some examples of Fano surfaces with supersingular reduction.

1. INTRODUCTION.

Let k be a finite field or a number field and let ℓ be a prime integer prime to the characteristic. Let $\bar{k}$ be an algebraic closure of k and let $G = \mathrm{Gal}(\bar{k}/k)$ be the Galois group. Let X be a smooth projective variety over k. We denote $\bar{X} = X \times_k \bar{k}$. Let $A^1(X)$ be the $\mathbb{Q}_\ell$-span in $H^2(\bar{X}, \mathbb{Q}_\ell(1))$ of the image of the divisor classes defined over k. The group G acts on $H^2(\bar{X}, \mathbb{Q}_\ell(1))$ and fixes the elements of $A^1(X)$. The Tate conjecture for divisors is as follows:

Conjecture 1. *(Tate Conjecture [29]). We have $A^1(X) = H^2(\bar{X}, \mathbb{Q}_\ell(1))^G$.*

When $k = \mathbb{F}_q$ is a finite field, the zeta function $Z(X,T)$ of X encodes the number of rational points of X over the extensions $k_r = \mathbb{F}_{q^r}$, and the Tate conjecture is equivalent to:

Conjecture 2. *([26], Proposition 4.3). The Picard number ρ_X of X is equal to the order of pole of the zeta function $Z(X, q^{-s})$ at $s = 1$.*

The Tate conjecture for abelian varieties has been proved by Tate for finite fields [27] and by Faltings for number fields [9]. Let k be a finite field or a number field, with characteristic $\neq 2$. Let $F \hookrightarrow \mathbb{P}^4_{/k}$ be a smooth cubic hypersurface. Through a generic point of $\bar{F}$ pass 6 lines of $\bar{F}$. The variety that parametrizes the lines on F is a smooth surface defined over k [1] called the Fano surface of lines of F. The surface S is minimal of general type and has invariants:

$$c_1^2 = 45,\ c_2 = 27,\ b_1 = 10,\ b_2 = 45.$$

We prove in this paper that:

Theorem 3. *The Tate conjecture holds for the surface S.*

This is a consequence of the following result:

Theorem 4. *Let S be a Fano surface defined over k. There is an isomorphism of Galois modules:*

$$\wedge^2 H^1(S,\mathbb{Q}_\ell) \simeq H^2(S,\mathbb{Q}_\ell).$$

For k a finite field, the Artin-Tate conjecture is a refinement of the Tate conjecture:

Conjecture 5. *(Artin-Tate Conjecture [28]). Let X be a smooth surface over k. The Brauer group $Br(X)$ of X is finite and:*

$$\lim_{s\to 1} \frac{P_2(X,q^{-s})}{(1-q^{1-s})^{\rho_X}} = \frac{(-1)^{\rho_X-1}|Br(X)|Disc(NS_k(X))}{q^{\alpha(X)}|NS_k(X)_{tor}|^2}$$

where $\alpha(X)=\chi(X,\mathcal{O}_X)-1+\dim \operatorname{Pic}^0(X)$, $NS_k(X)_{tor}$ is the torsion sub group of the Néron-Severi group $NS_k(X)$, and $Disc(NS_k(X))$ is the absolute value of the discriminant of $NS_k(X)/NS_k(X)_{tor}$ with respect to the intersection pairing.

By Milne [14], the Tate conjecture implies the Artin-Tate conjecture if the characteristic is not 2. In this paper, we give an algorithm that compute the zeta function $Z(S,T)$ of a Fano surface of a cubic F defined over finite fields of any characteristic $p>2$. We are therefore in position to compute the limit occurring in the Artin-Tate conjecture.

The intermediate Jacobian $J(F)$ of the cubic F is a 5-dimensional abelian variety canonically associated to F. This variety $J(F)$ is isomorphic to the Albanese variety of S and it is also a Prym variety $Pr(\Gamma,\mu)$ associated to a degree 2 étale cover μ of a certain plane quintic curve Γ. As we will see, it is equivalent to compute the zeta functions $Z(F,T), Z(S,T)$ or $Z(J(F),T)$.

In [12], Kedlaya gives an algorithm to compute the zeta function of cyclic cubic threefolds i.e., cubics that are triple cover of $\mathbb{P}^3$ branched over a smooth cubic surface. This algorithm runs by computing the number of points on the cubic hypersurface $F \hookrightarrow \mathbb{P}^4$ and uses the order 3 symmetries in order to reduce the computations. The algorithm we give here uses instead the Prym variety structure and computes the number of points on $J(F)$. It seems to be the first algorithm proposed in order to compute the zeta function of a Prym variety. We have implemented it in Sage (see [20]).

We end this paper by studying some interesting examples of cubic threefold and computing the zeta functions of some Fano surfaces over fields of characteristic $3,5$ and 7. Let ξ be a primitive (complex) 11^{th}-root of unity and let be $k=\mathbb{Q}(\xi)$. Let $S_{/k}$ be the Fano surface of lines of the Klein cubic threefold :

$$F_{/k}=\{x_1^2x_2+x_2^2x_3+x_3^2x_4+x_4^2x_5+x_5^2x_1=0\}.$$

The surface S has good reduction $S_\mathfrak{p}$ at every prime $\mathfrak{p}$ of k not dividing 11.

Corollary 6. *The Fano surface $S_{/k}$ has geometric Picard number 25.*
Let p be the prime dividing the norm of $\mathfrak{p}$. If p is odd and 11 is not a square modulo p, then $S_\mathfrak{p}$ is supersingular, i.e. its geometric Picard number equals $45=b_2$. If p is odd and 11 is a square modulo p, then $S_\mathfrak{p}$ has geometric Picard number 25.

To the knowledge of the author, there are only a few cases where the zeta function of surfaces have been computed : apart from the cases of abelian surfaces and a product of two curves, the zeta function has been computed for some examples of $K3$, some elliptic surface, for Fermat and some Delsarte hypersurfaces (see [24] and the references therein).

The Tate conjecture over finite fields is known for Fermat surfaces, products of curves, some K3, for abelian surfaces, surfaces with $p_g = 0$ with a lifting in characteristic 0, and surfaces X rationally dominated by a surface for which the Tate conjecture holds (see [14], [26] and the references). For examples of some Galois covers $S \to X$ where S is a Fano surface, see [21].

The Tate conjecture over number fields is known for K3, some Hilbert modular surfaces, many quaternionic Shimura surfaces and Picard surfaces (see [32], [26] and their references).

Acknowledgements. The author gratefully thanks Bert van Geemen for his suggestion to consider the conic bundle structure for computing the zeta function of cubics, and Kiran Kedlaya for emails exchanges. He thanks also Marc-Hubert Nicole and Stavros Papadakis for their comments on a previous version of this paper.
The author has been supported under FCT grant SFRH/BPD/72719/2010 and project Géometria Algébrica PTDC/MAT/099275/2008.

2. Notations, the trinity associated to a cubic threefold

The very rich geometry of cubic threefolds was a motivation for us to understand their arithmetic. These results are due to many authors, starting with Fano in the 40's and in historical order during the 70's: Bombieri, Swinnerton-Dyer [5], Clemens, Griffiths [7], Tyurin [30] [31], Murre [17] [18] [19], Altman, Kleiman [2] and Beauville [3]. There are other important papers on the subject since that time, but we will only recall the results we need.

Let F be a smooth cubic hypersurface defined over a field k and let S be its Fano surface. The Albanese variety A of S is 5 dimensional and defined over k. Let us suppose that S has a rational point s_o i.e. that F contains a line defined over k. Let $\vartheta : S \to A$ be the Albanese map such that $\vartheta s_o = 0$. Let Θ be the (reduced) image of $S \times S$ by the map $(s_1, s_2) \to \vartheta s_1 - \vartheta s_2$. The variety Θ is a divisor on A and (A, Θ) is a principally polarized abelian variety. Here we call the pair $J(F) = (A, \Theta)$ the intermediate Jacobian of F.

One of the most striking fact is that a Torelli-like theorem holds: for any intermediate Jacobian (A, Θ), we can recover (up to isomorphism) the original cubic F. This goes as follows : the divisor Θ has a unique singularity in 0. The exceptional divisor of the blow-up in 0 of A is a $\mathbb{P}^4$ and the closure of the strict transform of Θ into this $\mathbb{P}^4$ is (isomorphic to) the original cubic F.

Thus we see that given any one of the three items : the cubic F, its Fano surface S or its intermediate Jacobian (A, Θ), we can recover the two others. We can see therefore that there is one mathematical object which has three aspects, a trinity :

a) It is a cubic hypersurface F in $\mathbb{P}^4$ defined by one equation.
b) It is a smooth surface of general type S.
c) It is a 5-dimensional principally polarized abelian variety (A, Θ).

It seems interesting to us to compare this situation with the elliptic curves that are at the same time smooth cubic hypersurfaces of $\mathbb{P}^2$ and abelian varieties.

3. THE TATE CONJECTURE FOR FANO SURFACES.

Let A be an abelian variety of dimension ≥ 2 defined over a field k finitely generated over its prime field.

Definition 7. We say that a 2-dimensional effective cycle W on A is *non-degenerate* if the $\mathbb{Q}_\ell$-bilinear form:

$$Q_W : \begin{array}{ccc} H^2(\bar{A}, \mathbb{Q}_\ell(1)) \times H^2(\bar{A}, \mathbb{Q}_\ell(1)) & \to & H^{2n}(\bar{A}, \mathbb{Q}_\ell(n)) \simeq \mathbb{Q}_\ell \\ (x, y) & \to & x \cdot W \cdot y \end{array}$$

is non-degenerate, where we consider $W \in H^{2n-4}(\bar{A}, \mathbb{Q}_\ell(n-2))$.

For example, given an ample divisor Θ on A, the cycle $\frac{1}{(g-2)!}\Theta^{g-2}$ is positive by the Hodge-Riemann bilinear relations (see Chapter 0 Section 7 of [11]).

Let k be a finite field or a number field. Let S be a smooth surface over k with a k-rational point. Let A be the Albanese variety of S and let $\vartheta : S \to A$ be the Albanese map.

Proposition 8. *Suppose that the image W of S is a non-degenerate cycle and $b_2(S) = b_2(A)$. Then*

$$H^2(S, \mathbb{Q}_\ell) \simeq \wedge^2 H^1(S, \mathbb{Q}_\ell) \simeq \wedge^2 H^1(A, \mathbb{Q}_\ell) \simeq H^2(A, \mathbb{Q}_\ell).$$

The surface S satisfies the Tate conjecture and $\rho_S = \rho_A$.

We know only two examples which satisfy the conditions of the above Proposition:

Example 9. 1) For an abelian surface, the intersection pairing on $H^2(\bar{S}, \mathbb{Q}_\ell)$ is non-degenerate. Moreover: $H^2(S, \mathbb{Q}_\ell) = \wedge^2 H^1(S, \mathbb{Q}_\ell)$.
2) The Fano surface S of a smooth cubic threefold $F \hookrightarrow \mathbb{P}^4$ containing a rational line. The Albanese variety A of S is 5-dimensional and the Albanese map $\vartheta : S \to A$ is an embedding, moreover $b_2(S) = b_2(A) = 45$. There is a canonical principal polarisation Θ of A such that the class of S in A is $\frac{1}{3!}\Theta^3$ (see Clemens-Griffiths [7] and Tyurin [31]). The cycle S is therefore non-degenerate in A.

Proof. (Of Proposition 8). Let W be the image of S in the Albanese variety. For $x, y \in H^2(A, \mathbb{Q}_\ell(1))$, we have the following intersection numbers:

$$\vartheta^* x \vartheta^* y = (\deg \vartheta) x.W.y = (\deg \vartheta) Q_W(x, y),$$

and since Q_W is a non-degenerate pairing, the map

$$\vartheta^* : H^2(A, \mathbb{Q}_\ell(1)) \to H^2(S, \mathbb{Q}_\ell(1))$$

is injective. As $b_2(S) = b_2(A)$, the map ϑ^* is then an isomorphism of Galois modules $H^2(S,\mathbb{Q}_\ell) \simeq H^2(A,\mathbb{Q}_\ell)$. As the Tate conjecture is satisfied for divisors on abelian varieties over k (see [29] for finite fields and [9] for number fields), we have $\rho_A = \dim H^2(A,\mathbb{Q}_\ell(1))^G$. Since A is an abelian variety, we have

$$\wedge^2 H^1(A,\mathbb{Q}_\ell) \simeq H^2(A,\mathbb{Q}_\ell)$$

and since $H^1(S,\mathbb{Q}_\ell) \simeq H^1(A,\mathbb{Q}_\ell)$, we get

$$\wedge^2 H^1(S,\mathbb{Q}_\ell) \simeq \wedge^2 H^1(A,\mathbb{Q}_\ell) \simeq H^2(A,\mathbb{Q}_\ell) \simeq H^2(S,\mathbb{Q}_\ell).$$

Since ϑ^* is injective, we have $\rho_S \geq \rho_A$, moreover for every variety X, we have $\dim H^2(X,\mathbb{Q}_\ell(1))^G \geq \rho_X$ therefore:

$$\dim H^2(S,\mathbb{Q}_\ell(1))^G \geq \rho_S \geq \rho_A = \dim H^2(A,\mathbb{Q}_\ell(1))^G,$$

but as ϑ^* is an isomorphism $\dim H^2(S,\mathbb{Q}_\ell(1))^G = \dim H^2(A,\mathbb{Q}_\ell(1))^G$, thus we have equality$\rho_S = \rho_A$. □

4. The zeta function of a cubic threefold and its Fano surface.

4.1. **Notations, zeta functions of a variety.** Let X be a smooth n-dimensional variety over $k = \mathbb{F}_q$ such that $\bar{X}$ (the variety over an algebraic closure $\bar{k}$ of k) is smooth. Let N_r be the number of rational points of $\bar{X}$ over $\mathbb{F}_{q^r}$. The zeta function of X is defined by $Z(X,T) = exp(\sum_{r\geq 0} N_r \frac{T^r}{r})$ and can be written:

$$Z(X,T) = \sum_{i=0}^{i=2n} P_i(X,T)^{(-1)^{i+1}}$$

where $P_i(X,T)$ is the Weil polynomial with integer coefficients:

$$P_i(X,T) = \det(1 - T\pi^* | H^i(\bar{X},\mathbb{Q}_\ell)),$$

for π the Frobenius endomorphism. Suppose that X is a surface and let $Br(X)$ be the Brauer group of X. If the Tate conjecture is satisfied by X, then by Milne [14]:

Proposition 10. *(Artin-Tate Conjecture). Suppose the characteristic* > 2 *. The Brauer group of* X *is finite and:*

$$\lim_{s\to 1} \frac{P_2(X,q^{-s})}{(1-q^{1-s})^{\rho_X}} = \frac{(-1)^{\rho_X - 1}|Br(X)|Disc(NS_k(X))}{q^{\alpha(X)}|NS_k(X)_{tor}|^2}$$

where $\alpha(X) = \chi(X,\mathcal{O}_X) - 1 + \dim PicVar(X)$*,* $NS_k(X)_{tor}$ *is the torsion sub-group of* $NS_k(X)$*, and* $Disc(NS_k(X))$ *is the absolute value of the discriminant of* $NS_k(X)/NS_k(X)_{tor}$ *with respect to the intersection pairing.*

Let S be the Fano surface of a smooth cubic threefold $F \hookrightarrow \mathbb{P}^4$. The Picard variety of S is reduced [31] of dimension 5 and $\chi = 6$, therefore $\alpha(S) = 10$.

As $H^2(S,\mathbb{Q}_\ell) \simeq \wedge^2 H^1(A,\mathbb{Q}_\ell) \simeq H^2(A,\mathbb{Q}_\ell)$, the zeta function of S can be computed if we know the zeta function of A ; in fact we just need to know the action of the Frobenius on $H^1(A,\mathbb{Q}_\ell)$:

Theorem 11. *For characteristic* $\neq 2$*, we have:* $H^3(F,\mathbb{Q}_\ell) \simeq H^1(A,\mathbb{Q}_\ell(1))$.

Proof. By a result of Murre [17], Theorem 3 p. 149 asserting that $H^3(F,\mathbb{Q}_\ell) \simeq T_\ell(A) \otimes \mathbb{Q}_\ell$ where $T_\ell(A)$ is the Tate module of A. But $T_\ell(A)$ is isomorphic to $H^1(A,\mathbb{Q}_\ell)$. □

Remark 12. It would be interesting to have a bound on the order of $NS(S)_{tor}$. In the examples we give in Section 6, we compute the values of $\lim_{s\to 1} \frac{P_2(S,q^{-s})}{(1-q^{1-s})^{\rho_S}}$. For all the limits we computed, we get results on the form $\frac{N}{q^{10}}$, giving the idea that the group $NS(S)_{tor}$ should be trivial.

4.2. **Zeta function of a smooth cubic threefold.** The half upper part of the Hodge diamond of a smooth cubic threefold F over $\mathbb{C}$ is:

$$\begin{array}{ccccccc} & & & 1 & & & \\ & & 0 & & 0 & & \\ & 0 & & 1 & & 0 & \\ 0 & & 5 & & 5 & & 0 \end{array}$$

Thus $P_0(F,T) = 1-T$, $P_1(F,T) = 1 = P_5(F,T)$, $P_2(F,T) = 1-qT$. The polynomials P_4 and P_6 are computed by Poincaré duality: $P_4 = 1 - q^2T$, $P_6(F,T) = 1-q^3T$.

Corollary 13. *We have:*

$$Z(F,T) = \frac{P_3(F,T)}{(1-T)(1-qT)(1-q^2T)(1-q^3T)},$$

and

$$N_r(F) = (1+q^r+q^{2r}+q^{3r}) - q^r Tr(\pi^r|H^1(\bar{A},\mathbb{Q}_\ell)),$$

where π is the Frobenius endomorphism of $\bar{A}$.

The roots of $P_i(F,T)$ have absolute value $q^{-i/2}$. We have:

$$P_1(S,T) = P_1(A,T) = P_3(F,\frac{T}{q}).$$

Since by a computer it is possible to compute the number of points of F, it is possible to get the zeta function of F, A and S. Kedlaya's algorithm compute the number of points of F. Alternatively, we can also compute the number of points of A that is a Prym variety of a degree 2 étale cover of a certain plane quintic, this is the idea of the algorithm we propose.

4.3. **Zeta function and Picard number of a Fano surface.** Let S be the Fano surface of a smooth cubic $F \hookrightarrow \mathbb{P}^4$ over $\mathbb{F}_q$. We have

$$Z(S,T) = \frac{P_1(S,T)P_3(S,T)}{(1-T)P_2(S,T)(1-q^2T)}.$$

We can write $P_1 = \prod_{i=1}^{i=10}(1-\omega_i T)$ with $|\omega_i| = q^{-1/2}$. As

$$H^2(S,\mathbb{Q}_\ell) = \wedge^2 H^1(S,\mathbb{Q}_\ell)$$

we get $P_2(S,T) = \prod_{1\leq i<j\leq 10}(1-\omega_i\omega_j T)$. Moreover $P_3(S,T) = q^{15}T^{10}P_1(S,\frac{1}{q^2T})$. Thus

$$Z(S,T) = \frac{\prod_{i=1}^{i=10}(1-\omega_i T)\prod_{i=1}^{i=10}(1-\frac{q^2}{\omega_i}T)}{(1-T)(1-q^2T)\prod_{1\leq i<j\leq 10}(1-\omega_i\omega_j T)}.$$

The order of the pole at 1 of $Z(S,q^{-s})$ equals the number of elements of the set $\{(i,j)/1 \leq i < j \leq 10 \text{ and } \omega_i\omega_j = q\}$; it is the multiplicity of the root $1/q$ in $P_2(S,T) = P_2(A,T)$. By the Tate conjecture for abelian varieties, we get:

Corollary 14. *The Fano surface and the intermediate Jacobian have Picard numbers at least* 5*.*

Let $N_r(X)$ be the number of rational points of a variety X over $\mathbb{F}_{q^r}$. For each Weil polynomial $P_i(X,T)$, let $\omega_{i,1},\dots,\omega_{i,b_i}$ be the reciprocal roots of P_i. The formula expressing the zeta function gives:

$$N_r(X) = \sum_{i=0}^{i=2\dim X} (-1)^i \sum_{j=1}^{j=b_i} \omega_{i,j}^r.$$

We will use this formula for computing the numbers $N_1(S), N_2(S)$. For an abelian variety A, we have $N_1(A) = P_1(A,1)$.

We will need also the following Theorem of Honda-Tate:

Theorem 15. *(*[16]*, Theorems 2-3, Appendix I or* [8]*). Let A and B be abelian varieties over a finite field k of q elements. Let $f_A, f_B \in \mathbb{Z}[T]$ be the characteristic polynomials of their q-Frobenius endomorphisms, so f_A and f_B have degrees $2\dim A$ and $2\dim B$ with constant terms $q^{\dim A}$ and $q^{\dim B}$ respectively. The following statements are equivalent:*
(1) B is k-isogenous to an abelian subvariety of A.
(2) f_A divides f_B in $\mathbb{Q}[T]$.
In particular, A is k-isogenous to B if and only if $f_A = f_B$. Moreover, A is isogenous to a power of one k-simple abelian variety if and only if f_A is a power of an irreducible polynomial in $\mathbb{Q}[T]$.

5. The intermediate Jacobian as a Prym variety.

Let $F \hookrightarrow \mathbb{P}^4$ be a smooth cubic threefold defined over a field k and let L be a line on F defined over k. Let us denote by C_L the incidence curve parametrizing the lines on F cutting L. Let Γ be the curve parametrizing planes Y containing L and such that the degree 3 plane curve $Y \cdot F$ is the union of three lines. The curve C_L has (arithmetic) genus 11 and Γ is a plane quintic of arithmetic genus 6.

Theorem 16. *(Murre* [19]*) For generic L, the curve C_L is smooth and there is a degree 2 étale map $C_L \to \Gamma$ such that the associated Prym variety $Pr(C_L)$ is isomorphic to the Albanese variety of the Fano surface S.*

Let us suppose that $k = \mathbb{F}_q$ is the finite field with q elements, of characteristic not 2. The aim of this section is to give an algorithm that compute the zeta function of S and the Albanese variety A by using the Prym variety structure. We use here the reasoning of [5].

Let $x_1,\dots,x_5$ be coordinates in $\mathbb{P}^4$ and let $L \hookrightarrow F$ be a k-rational line on F. We can suppose that $L = \{x_1 = x_2 = x_3 = 0\}$ and $F = \{F_{eq} = 0\}$ with

$$F_{eq} = \ell_1 x_4^2 + 2\ell_2 x_4 x_5 + \ell_3 x_5^2 + 2q_1 x_4 + 2q_2 x_5 + f,$$

where $\ell_1, \ell_2, \ell_3, q_1, q_2, f$ are homogenous forms in x_1, x_2, x_3 of respective degree $1,1,1,2,2,3$. Let $X \hookrightarrow \mathbb{P}^4$ be the plane $x_4 = x_5 = 0$. Any plane $Y \hookrightarrow \mathbb{P}^4$ containing the line L cuts X into a unique point. The plane X is thus the variety of planes in $\mathbb{P}^4$ containing L. For a $\bar{k}$-point $x = (x_1 : x_2 : x_3 : 0 : 0)$ of $X = \mathbb{P}^2$, we denote by Y_x the unique plane containing x. The intersection of F and Y_x has equation $F_{eq}(y_1x_1 : y_1x_2 : y_1x_3 : y_2 : y_3) = 0$, i.e.:

$$y_1(\ell_1 y_2^2 + 2\ell_2 y_2 y_3 + \ell_3 y_3^2 + 2q_1 y_1 y_2 + 2q_2 y_1 y_3 + y_1^2 f) = 0$$

in the plane Y_x with coordinates y_1, y_2, y_3. Therefore, the intersection $Y \cdot F$ equals $L + Q$ where $Q = Q_x$ is the residual quadric with equation:

$$Q = \{\ell_1 y_2^2 + 2\ell_2 y_2 y_3 + \ell_3 y_3^2 + 2q_1 y_1 y_2 + 2q_2 y_1 y_3 + y_1^2 f = 0\}.$$

The quadric Q_x is not irreducible over $\bar{k}$ if and only if Q_x is singular. The scheme parametrizing the planes Y_x such that Q_x decomposes (over an extension) as $Q = L_1 + L_2$ with L_1, L_2 two lines, is a plane reduced quintic curve $\Gamma \hookrightarrow X = \mathbb{P}^2(x_1, x_2, x_3)$ defined over k, of equation $\det M_\Theta = 0$, where:

$$M_\Theta = \begin{pmatrix} \ell_1 & \ell_2 & q_1 \\ \ell_2 & \ell_3 & q_2 \\ q_1 & q_2 & f \end{pmatrix}.$$

Let C_L be the incidence scheme parametrizing the lines that cut L.

Lemma 17. [5] *The curve Γ has at most nodal singularities. The curve Γ is singular at the points x such that $Y_x \cdot F = L + 2L'$, where L' is a line. The curve Γ is smooth for L generic.*
The scheme C_L is a reduced one dimensional subscheme of the Fano surface S. The is a natural degree 2 map $\mu : C_L \to \Gamma$ that is ramified precisely over the singularities of Γ. The point p of C_L is smooth if and only if $\mu(p)$ is smooth.

Let $\tilde{F} \to F$ be the blow-up of F along the line L. The variety $\tilde{F}$ has a natural structure of a conic bundle ([4]):

$$\tilde{F} \to \mathbb{P}^2 = X$$

where $\mathbb{P}^2$ is the Grassmanian of planes containing the line L. The fiber over the point x is (isomorphic to) the quadric $Q = Q_x$ such that $F \cdot Y_x = Q + L$. For x defined over k, the number of k-points of Q_x is:
i) $q + 1$ if Q is geometrically irreducible i.e. $x \notin \Gamma(k)$ or if $x \in \Gamma(k)$ is a singular point of Γ, in that case $Q = 2L'$ with L' a k-rational line on F.
ii) $2q + 1$ if $Q = L_1 + L_2$ with L_1, L_2 two different lines defined over k, i.e. the 2 points in C_L over $x \in \Gamma(k)$ are in $C_L(k)$.
iii) 1 if Q degenerates to a union of two line over a degree 2 extension of k, i.e. if the 2 points in C_L over $x \in \Gamma(k)$ are not in $C_L(k)$.

Let $N_r(X)$ denotes the number of rational points of a variety X over the degree r extension $k_r = \mathbb{F}_{q^r}$ of k.

Proposition 18. *([5], formula (18)). We have*

$$N_r(F) = q^{3r} + q^{2r} + q^r + 1 + q^r(N_r(C_L) - N_r(\Gamma)).$$

This is proved in [5], however we give their proof here because it explains the algorithm computing $N_r(F)$ we describe below.

Proof. Let a be the number of k-rational singularities of Γ (and C_L). Taking care of the three above possibilities i), ii) and iii), we get:

$$\begin{aligned} N_1(\tilde{F}) = \quad & (q+1)(N_1(\mathbb{P}^2) - N_1(\Gamma) + a) + (2q+1)\tfrac{1}{2}(N_1(C_L) - a) \\ & +(N_1(\Gamma) - a - \tfrac{1}{2}(N_1(C_L) - a)) \end{aligned}$$

thus

$$N_1(\tilde{F}) = q^3 + 2q^2 + 2q + 1 + q(N_1(C_L) - N_1(\Gamma)).$$

As each point on $L \hookrightarrow F$ is replaced by a $\mathbb{P}^1$ on $\tilde{F}$, we have moreover:

$$N_1(\tilde{F}) = N_1(F) - (q+1) + (q+1)^2,$$

thus

$$N_1(F) = q^3 + q^2 + q + 1 + q(N_1(C_L) - N_1(\Gamma)),$$

and this proves Proposition 18. □

Let $\mu : C_L \to \Gamma$ be the degree 2 map and let x be a k-rational smooth point. In order to compute the numbers $N_r(C_L) - N_r(\Gamma)$, we need to understand when the 2 $\bar{k}$-rational points in $\mu^* x$ are k-rational. For $1 \leq i \leq 3$, let δ_i be the determinant of the order 1 (i,i)-minor of the matrix M_Θ. We consider δ_i as a section of $H^0(\Gamma, \mathcal{O}_\Gamma(1))$.

Proposition 19. *For every smooth point x of Γ there exists an integer $1 \leq i = i(x) \leq 3$ such that $\delta_i(x) \neq 0$. The curve C_L has two rational points over $x \in \Gamma(k)$ if and only if $-\delta_i(x) \in (k^*)^2$.*

Proof. Let x a k-rational smooth point of Γ and let $Q = Q_x$ such that $F \cdot Y_x = L + Q$. The line $L = \{y_1 = 0\}$ cuts $Q \hookrightarrow Y_x$ in the points such that:

$$y_1 = \ell_1 y_2^2 + 2\ell_2 y_2 y_3 + \ell_3 y_3^2 = 0.$$

Therefore, if $-\delta_3(x) = \ell_2^2(x) - \ell_1(x)\ell_2(x)$ is nonzero, the curve C_L has two rational points over $x \in \Gamma(k)$ if and only if $-\delta_3(x) \in (k^*)^2$.
If $\delta_3(x) = 0$, we only sketch the proof and refer to [4], Lemme 1.6 and its proof. In that case, we have $Y_x \cdot F = L + Q_x = L + L_1 + L_2$ with L_1, L_2 defined over $\bar{k}$ and meeting in a k-rational point p of L. It is possible to explicitly compute a model in $\mathbb{P}^3$ of (the degree 6) scheme $\mathcal{X}_p$ of lines going through p. By knowing $\mathcal{X}_p$, we can determine whether the two points on $\mathcal{X}_p$ corresponding to the lines L_1, L_2 are k-rational or not and this is so if and only if $-\delta_1$ is a nonzero square or $-\delta_2$ is a nonzero square. □

Let us describe the algorithm for the computation of the numbers $N_r(F)$ and $N_r(C_L) - N_r(\Gamma)$.

The input data is a cubic threefold F over $\mathbb{F}_q$ containing a $\mathbb{F}_q$-rational line L. To this data is associated the matrix

$$M_\Theta = \begin{pmatrix} \ell_1 & \ell_2 & q_1 \\ \ell_2 & \ell_3 & q_2 \\ q_1 & q_2 & f \end{pmatrix}$$

whose determinant is the equation of the quintic $\Gamma \hookrightarrow X = \mathbb{P}^2$ (maybe singular). Then we compute $N_r = N_r(C_L) - N_r(\Gamma)$ as follows:

Initiate $N_r := 0$. For $x \in \mathbb{P}^2(\mathbb{F}_{q^r})$, if $\det(M_\Theta)(x) = 0$ then if $-\delta_3(x) \in (\mathbb{F}_{q^r}^*)^2$ then $N_r := N_r + 1$, else if $-\delta_3(x) \neq 0$, then $N_r := N_r - 1$, otherwise if $-\delta_1(x) \in (\mathbb{F}_{q^r}^*)^2$ then $N_r := N_r + 1$ else if $-\delta_1(x) \neq 0$, then $N_r := N_r - 1$, otherwise if $-\delta_2(x) \in (\mathbb{F}_{q^r}^*)^2$ then $N_r := N_r + 1$ else if $-\delta_2(x) \neq 0$ then $N_r := N_r - 1$ end if, end for.

The output N_r equals $N_r(C_L) - N_r(\Gamma)$. Remark that the $-\delta_i$ are transition functions of an invertible sheaf $\mathcal{L}$ on Γ such that $\mathcal{L}^{\otimes 2} = \omega_\Gamma$ that corresponds to the degree 2 cover $\mu : C_L \to \Gamma$. Thus a point on Γ is singular if and only if $\forall 1 \leq i \leq 3$, $\delta_i(x) = 0$.

The knowledge of $N_1, \dots, N_5$ is thus enough to get the 5 first coefficients of $P_1(Pr((\Gamma, \mu), T)$ and the remaining 5 ones are determined by the symmetries.

Remark 20. (1) This algorithm for computing the action of the Frobenius on the Prym variety $Pr(C_L)$ is generalizable to other plane curves occurring as discriminant locus of some quadric bundles, see [4].
(2) If the characteristic of k equals 2, by [5] end of page 6, it is possible to adapt the algorithm computing $N_r(F)$ if the line L is chosen sufficiently generic such that Γ is smooth (see also [4], remarque 2).

6. EXAMPLES

6.1. **Reduction in characteristic** $3, 5$ **and** 7 **of a generic cubic.** Let $F \hookrightarrow \mathbb{P}^4$ be the cubic threefold with equation:

$$F_{eq} = x_1 x_4^2 + 2x_2 x_4 x_5 + x_3 x_5^2 + 2q_1 x_4 + 2q_2 x_5 + f,$$

where:

$$\begin{gathered} q_1 = x_1^2 + 2x_2^2 + x_2 x_3 + x_3^2 \\ q_2 = x_1 x_2 + 4x_2 x_3 + x_3^2 \\ f = x_2^2 x_3 - (x_1^3 + 4x_1 x_2^2 + 2x_2^3). \end{gathered}$$

The cubic F is smooth in characteristic $5, 7, 11$ and 13 and singular in characteristic $2, 3$. The associated quintic curve Γ is smooth in characteristic $3, 7, 11, 13$, but singular in characteristic $2, 5$.

Remark 21. This example is interesting because the cubic $F_{/\mathbb{Q}}$ and its Fano surface $S_{/\mathbb{Q}}$ have bad reduction at the place 3, however the intermediate Jacobian $J(F)$ has good reduction. This is the same phenomena as for the curves and their Jacobian.
Moreover, in characteristic 0, Casalaina-Martin and Friedman [6] proved that the cubic associated to a plane quintic curve provided with a theta characteristic M_Θ is always smooth. This example over $\mathbb{F}_3$ is a counter example. However, by looking to [6], this phenomena seems to hold only in characteristic 3.
Lastly, we remark that the curves Γ and C_L both have bad reduction at the place 5, but the associated Prym variety has good reduction.

We have implemented the algorithm in Sage [20]. Using a personal laptop, it takes 5 minutes in characteristic 7 to obtain $P_1(S, T)$.

Over $\mathbb{F}_3$, as the cubic $F_{/\mathbb{F}_3}$ is singular, we can only compute the Weil polynomial $P_1(Pr(\Gamma,\mu)_{/\mathbb{F}_3},T)$ where $\mu : C_L \to \Gamma$ is the étale double cover associated to the quintic Γ and its theta characteristic M_Θ. We get:

$$\begin{aligned} P_1(Pr(\Gamma,\mu)_{/\mathbb{F}_3},T) = \ & 1+3T+11T^2+22T^3+49T^4+79T^5+147T^6 \\ & +198T^7+297T^8+243T^9+243T^{10}. \end{aligned}$$

The Prym variety $Pr(\Gamma,\mu)_{/\mathbb{F}_3}$ has geometric Picard number 5.

Over $\mathbb{F}_5$, we get:

$$P_1(S_{/\mathbb{F}_5},T) = (5T^2+1)(625T^8+50T^6+40T^5-6T^4+8T^3+2T^2+1).$$

The Fano surface $S_{/\mathbb{F}_5}$ has Picard number 5 and has 33 $\mathbb{F}_3$-rational points. We have:

$$\lim_{s\to 1} \frac{P_2(S_{/\mathbb{F}_5},5^{-s})}{(1-5^{1-s})^5} = \frac{2^{18}\cdot 3^5 \cdot 157}{5^{10}}.$$

Over $\mathbb{F}_7$, we get:

$$\begin{aligned} P_1(S_{/\mathbb{F}_7},T) = \ & 1+4T+15T^2+46T^3+159T^4+460T^5+1113T^6 \\ & +2254T^7+5145T^8+9604T^9+16807T^{10}. \end{aligned}$$

It is an irreducible polynomial over $\mathbb{Q}$, therefore by the Honda-Tate Theorem, the intermediate jacobian $J(F)$ of F is simple. The Fano surface $S_{/\mathbb{F}_7}$ has Picard number 5 and 97 $\mathbb{F}_7$-rational points. We have:

$$\lim_{s\to 1} \frac{P_2(S_{/\mathbb{F}_7},7^{-s})}{(1-7^{1-s})^5} = \frac{2^4\cdot 83^2 \cdot 557 \cdot 5737}{7^{10}}.$$

Remark 22. We give a proof in [23] that a generic Fano surface over $\mathbb{C}$ has Picard number $\rho = 1$. One would like to exhibit an example of a Fano surface over $\mathbb{Q}$ with $\rho = 1$. However, by reducing the above Fano surface $S_{/\mathbb{Q}}$ modulo a prime gives us only the bound $\rho \leq 5$.

Over the field $\mathbb{F}_{11}$, the computation becomes impracticable : it needs 4 minutes to get N_4 but more that 24 hours to get N_5. By [12] Lemma 1.2.3, for positive integers q,d,j and complex numbers $a_1,\dots,a_{j-1}$, there exist a certain disk of radius $\frac{d}{j}q^{j/2}$ which contains every a_j for which we can choose $a_{j+1},\dots,a_d \in \mathbb{C}$ so that the polynomial

$$R(T) = 1 + \sum_{j=1}^{j=d} a_j T^j$$

has all roots on the circle $|T| = q^{-1/2}$. In our case, $\frac{d}{j}q^{j/2} = \frac{10}{5}11^{5/2} = 802.623...$ and we get $P_1(S_{/\mathbb{F}_{11}},T) = P_a$, where a is an integer in $\{80,\dots,332\}$ and

$$\begin{aligned} P_a = \ & 1-T+13T^2+T^3-28T^4+aT^5-11\cdot 28T^6 \\ & +11^2T^7+11^3\cdot 13T^8-11^4T^9+11^5T^{10} \end{aligned}$$

For all the consecutive values $a \in \{80,\dots,332\}$, the polynomial P_a has its roots equal to $11^{-1/2}$ (with error at most 10^{-10}) and we have not been able to find the a corresponding to the Fano surface. The same problem holds over $\mathbb{F}_{13}$ for 63 consecutive values.

6.2. **Fano surfaces with 12 elliptic curves.** Let K be a number field and let $\lambda \in K$. Let us recall:

Proposition 23. *([22]) The Albanese variety of the Fano surface of the cubic*

$$F_\lambda = \{x_1^3 + x_2^3 + x_3^3 + \lambda x_1x_2x_3 + x_4^3 + x_5^3\} \hookrightarrow \mathbb{P}^4_{/K}$$

is isogenous (over $\bar{K}$) to $E_0^3 \times E_\lambda^2$, where E_λ is the elliptic curve $x_1^3 + x_2^3 + x_3^3 - \lambda x_1x_2x_3 = 0$.

If E_λ is not isomorphic to E_0, the Fano surface of F contains 12 elliptic curves ([22]). Let be α a primitive third root of unity. The curve E_0 has complex multiplication by $\mathbb{Z}[\alpha]$. Let us choose $\lambda \in K$ such that E_λ has complex multiplication by an order of $\mathbb{Z}[\alpha]$. Let $p > 2$ be a prime and let $\mathfrak{p}$ be a prime in K over p. Suppose that F_λ, S and $J(F_\lambda)$ have good reduction at $\mathfrak{p}$ and let $S_\mathfrak{p}$ be the reduction at $\mathfrak{p}$ of S.

Proposition. *If $p = 2$ mod 3, the surface $S_\mathfrak{p}$ is supersingular : its geometric Picard number equals $b_2(S) = 45$.*

Proof. By the choice of λ, the intermediate jacobian $J(F)$ is isogenous over $\bar{K}$ to E_0^5. As for p prime with $p = 2$ mod 3 the curve E_0 has supersingular reduction, the geometric Picard number of the reduction mod $\mathfrak{p}$ of $J(F)$ equals 45 (see [16]). □

Let us consider the cubic

$$F_\lambda = \{x_1^3 + x_2^3 + x_3^3 + \lambda x_1x_2x_3 + x_4^3 + x_5^3\} \hookrightarrow \mathbb{P}^4_{/\mathbb{F}_7}$$

over $\mathbb{F}_7$. For $\lambda = 3$, we get by Kedlaya's algorithm:

$$P_1(S,T) = (1 + T + 7T^2)^2(1 + 4T + 7T^2)^3.$$

The intermediate Jacobian of F_λ has Picard number 13. We have:

$$\lim_{s\to 1} \frac{P_2(S, 7^{-s})}{(1 - 7^{1-s})^{13}} = \frac{2^6 \cdot 3^{18}}{7^{10}}.$$

Moreover:

$$\lim_{s\to 1} \frac{P_2(S/\mathbb{F}_{49}, 49^{-s})}{(1 - 49^{1-s})^{25}} = \frac{2^{18} \cdot 3^{18} \cdot 5^{12}}{7^{20}}.$$

The Fano surface has 270 rational points over $\mathbb{F}_7$.

6.3. **The Klein cubic threefold.** Let ξ be a primitive (complex) root of unity and let be $K = \mathbb{Q}(\xi)$. Let $S_{/K}$ be the Fano surface of lines of the Klein cubic threefold :

$$F_{/K} = \{x_1^2x_2 + x_2^2x_3 + x_3^2x_4 + x_4^2x_5 + x_5^2x_1 = 0\}.$$

It is easy to check that S has good reduction $S_\mathfrak{p}$ at every prime $\mathfrak{p}$ not over 11. Let p be the prime dividing the norm of $\mathfrak{p}$.

Proposition 24. *If p is odd and 11 is not a square modulo p, then $S_\mathfrak{p}$ is supersingular i.e. its geometric Picard number equals $45 = b_2$. If p is odd and 11 is a square modulo p, then $S_\mathfrak{p}$ has geometric Picard number 25.*

Proof. Let be $\nu = \frac{-1+\sqrt{-11}}{2}$ and $E = \mathbb{C}/\mathbb{Z}[\nu]$. In [23], we prove that $J(F)_{/\mathbb{C}}$ is isomorphic to E^5. By [25], App. A3, the elliptic curve E has the following model over $\mathbb{Q}$:

$$y^2 + y = x^3 - x^2 - 7x + 10,$$

which we still denote by E. The curve E has good reduction for prime $p \neq 11$. Let be $\mathcal{O} = \mathbb{Z}[\nu]$, then the curve E has complex multiplication by $\mathcal{O}$ (over a certain extension). We use the criteria of Deuring ([13] Chap 13, Thm 12) : for odd $p \neq 11$, the reduction of E modulo p is a supersingular if and only if p is inert or ramified in $\mathcal{O}$. By classical results on number theory, an odd prime $p \neq 11$ is inert or ramified in $\mathcal{O}$ if and only if 11 is not a square modulo p.

Over an extension, the intermediate Jacobian $J(F)$ is isogenous to E^5. By [16], the geometric Picard number of the reduction modulo $\mathfrak{p}$ of $J(F)$ is therefore 45 if 11 is not a square modulo p, and 25 otherwise. □

References

[1] Achter J., "Arithmetic Torelli maps for cubic surfaces and threefolds", preprint arXiv:1005.2131v3

[2] Altman A., Kleiman S., "Foundations of the theory of Fano schemes", Compositio Math. 34 (1977), no. 1, 3–47.

[3] Beauville A., "Les singularités du diviseur Θ de la jacobienne intermédiaire de l'hypersurface cubique dans $\mathbb{P}^4$", Algebraic threefolds (Varenna, 1981), pp. 190–208, Lecture Notes in Math., 947, Springer, Berlin-New York, 1982.

[4] Beauville A., "Variétés de Prym et Jacobiennes intermédiaires", Ann. Scient. Éc. Norm. Sup. 10 (1977), 309–391.

[5] Bombieri E., Swinnerton-Dyer H.P.F., "On the local zeta function of a cubic threefold", Ann. Scuola Norm. Sup. Pisa (3) 21 1967, 1–29.

[6] Casalaina-Martin S., Friedman R., "Cubic threefolds and abelian varieties of dimension five", J. Algebraic Geom. 14 (2005), no. 2, 295–326.

[7] Clemens H., Griffiths P., "The intermediate Jacobian of the cubic threefold", Ann. of Math. (2) 95 (1972), 281–356

[8] Eisenträger K., "The Theorem of Honda and Tate", notes available at http://www.math.psu.edu/eisentra/hondatate.pdf

[9] Faltings G., "Endlichkeitssätze für abelsche Varietäten über Zahlkörpern", Invent. Math. 73 (1983), no. 3, 349–366.

[10] Fano G., "Irrazionalità della forma cubica generale dello spazio a quattro dimensioni", Univ. e Politecnico Torino. Rend. Sem. Mat. 9, (1950). 21–32.

[11] Griffiths P., Harris J., "Principles of algebraic geometry", Wiley Classics Library. John Wiley and Sons, Inc., New York, 1994. xiv+813 pp

[12] Kedlaya K., "Effective p-adic cohomology for cyclic cubic threefolds", to appear in Computational Algebraic and Analytic Geometry of Low-dimensional Varieties, Contemporary Mathematics, AMS.

[13] Lang S., "Elliptic Functions", Addison-Wesley Publishing Co., Inc., Reading, Mass.-London-Amsterdam, 1973. xii+326 pp.

[14] Milne J., "On a conjecture of Artin and Tate", Ann. of Math. (2) 102 (1975), no. 3, 517–533.

[15] Mumford D., "Red book of Varieties and Schemes", Second, expanded edition. Includes the Michigan lectures (1974) on curves and their Jacobians. With contributions by Enrico Arbarello. Lecture Notes in Mathematics, 1358. Springer-Verlag, Berlin, 1999. x+306 pp.

[16] Mumford D., "abelian Varieties", Tata Institute of Fundamental Research Studies in Mathematics, No. 5 Published for the Tata Institute of Fundamental Research, Bombay; Oxford University Press, London, 1970.

[17] Murre J.P., "Some results on cubic threefolds", Classification of algebraic varieties and compact complex manifolds, pp. 140–160. Lecture Notes in Math., Vol. 412, Springer, Berlin, 1974.

[18] Murre J. P., "Reduction of the proof of the non-rationality of a non-singular cubic threefold to a result of Mumford", Compositio Math. 27 (1973), 63–82.

[19] Murre J. P., "Algebraic equivalence modulo rational equivalence on a cubic threefold", Compositio Math. 25 (1972), 161–206.

[20] Roulleau X., Sage and Maple worksheets available in http://web.me.com/xavier.roulleau/ (publications).

[21] Roulleau X., "Quotients of Fano surfaces", to appear in Rend. Lincei.

[22] Roulleau X., "Fano surfaces with 12 or 30 elliptic curves", Michigan Math. J. 60 (2011), no. 2, 313–329

[23] Roulleau X.,"The Fano surface of the Klein cubic", J. Math. Kyoto Univ. 49 (2009), no. 1, 113–129.

[24] Schuett M., "K3 surfaces with non-symplectic automorphisms of 2-power order", J. Algebra 323 (2010), no. 1, 206–223.

[25] Silverman J. "Advanced topics in the arithmetic of Elliptic curves", Graduate Texts in Mathematics, 151. Springer-Verlag, New York, 1994.

[26] Tate J., "Conjectures on Algebraic cycles in ℓ-adic cohomology", Motives (Seattle, WA, 1991), 71–83, Proc. Sympos. Pure Math., 55, Part 1, Amer. Math. Soc., Providence, RI, 1994

[27] Tate, J., "Endomorphisms of abelian varieties over finite fields", Invent. Math. 2 1966 134–144.

[28] Tate J., "On a conjecture of Birch and Swinnerton-Dyer and a geometric analog", Séminaire Bourbaki, 1964-1966, exp. no 306, p. 415–440.

[29] Tate J., "Algebraic cycles and poles of zeta functions", 965 Arithmetical Algebraic Geometry (Proc. Conf. Purdue Univ., 1963) pp. 93–110 Harper and Row, New York

[30] Tyurin, A.N., "The geometry of the Fano surface of a nonsingular cubic $F \subset \mathbb{P}^4$ and Torelli Theorems for Fano surfaces and cubics", Math. Ussr Izv. 5, 517–546 (1971)

[31] Tyurin A.N., "The Fano surface of a nonsingular cubic in $\mathbb{P}^4$", Izv. Akad. Nauk SSSR Ser. Mat. 34 1970 1200–1208.

[32] Virdol C., "Tate conjecture for twisted Picard modular surfaces", J. Number Theory 131 (2011), no. 6, 1048–1053.

Xavier Roulleau,
`roulleau@math.ist.utl.pt`
Départemento de Mathematica,
Instituto Superior Técnico,
Avenida Rovisco Pais
1049-001 Lisboa
Portugal